

# Physics 207 – Lecture 23

**Physics 207, Lecture 28, Dec. 11**

- Agenda: Ch. 21 Finish, Start Ch. 22
  - ❖ Ideal gas at the molecular level, Internal Energy
  - ❖ Molar Specific Heat ( $Q = m c \Delta T = n C \Delta T$ )
  - ❖ Ideal Gas Molar Heat Capacity (and  $\Delta U_{int} = Q + W$ )
    - Constant V:  $C_v = 3/2 R$ , Constant P:  $C_p = 3/2 R + R = 5/2 R$
  - ❖ Adiabatic processes (no heat transfer)
  - ❖ Heat engines and Second Law of thermodynamics
  - ❖ Reversible/irreversible processes and Entropy


Assignments:

- Problem Set 10 (Ch. 20 & 21) due Tuesday, Dec. 12, 11:59 PM  
Ch. 20: 13,22,38,43,50,68 Ch.21: 2,16,29,36,70
- Problem Set 11, Ch. 22: 6, 7, 17, 37, 46 (Due, Friday, Dec. 15, 11:59 PM)
- Wednesday, Work on problem set 11

Physics 207: Lecture 28, Pg 1

**Lecture 28, Exercise 1**

- An atom in a classical solid can be characterized by three independent harmonic oscillators, one for the x, y and z-directions?
- How many degrees of freedom are there?



(A) 1 (B) 2 (C) 3 (D) 4 (E) Some other number

Physics 207: Lecture 28, Pg 2

**Ideal Gas Molar Heat Capacities**

- Definition of molar heat capacities (relates change in the internal energy to the change in temperature)

Ideal Gas Internal Energy

$$C \equiv \frac{1}{n} \lim_{n \rightarrow \infty} Q / \Delta T = \frac{1}{n} \delta Q / \delta T \quad K_{tot trans} = U = \frac{3}{2} N k_B T = \frac{3}{2} n R T$$


- There is only microscopic kinetic energy (i.e., no springs) in a monoatomic ideal gas (He, Ne, etc.)
  - ❖ At constant V: Work W is 0 so  $\Delta U = Q \rightarrow C_v = \frac{3}{2} R$
  - ❖ At constant P:  $\Delta U = Q + W = Q - P \Delta V$   $C_p = \frac{3}{2} R + R$

$PV = nRT$

Physics 207: Lecture 28, Pg 3

**Lecture 28, Exercise 2**

- An atom in a classical solid can be characterized by three independent harmonic oscillators, one for the x, y and z-directions ( U per atom = 3  $k_B T$  ) ?
- What is the classical molar heat capacity (P  $\Delta V \neq 0$  !)?



(A) nR (B) 2nR (C) 3nR (D) 4nR (E) Some other number

Physics 207: Lecture 28, Pg 4

**Adiabatic Processes**

- By definition a process in which no heat transfer (Q) occurs


For an Ideal Gas:

$$PV^\gamma = \text{const} \quad \gamma \equiv \frac{C_p}{C_v}$$

- Adiabatic process:
  - ❖  $PV^\gamma$  is constant
  - ❖  $PV = nRT$  but **not** isothermal
  - ❖ Work (on system) becomes :

$$W = -\int_{V_1}^{V_2} P dV = -\int_{V_1}^{V_2} \frac{\text{const}}{V^\gamma} dV = \frac{\text{const}}{\gamma} (V_2^{-\gamma} - V_1^{-\gamma})$$

Physics 207: Lecture 28, Pg 5



### Evaporative Cooling in a Bose-Einstein Condensation

Evaporative cooling can lead to a state change

Physics 207: Lecture 28, Pg 7

### Granularity, Energy and the Boltzmann Statistics

- There are discrete number accessible energy levels in any finite system. It can be shown that if there are many more levels than particles to fill them the probability is just
 
$$P(E) = \exp(-E/k_B T)$$
- The energy levels for a Quantum Mechanical (i.e., discrete quantized states) ideal gas is shown before and after a change (highly idealized diagram, imagine lots more levels and lots more particles).
- Here we increase the box size slowly and perform a "quasistatic, adiabatic expansion"

Physics 207: Lecture 28, Pg 8

### Chapter 22: Heat Engines and the 2<sup>nd</sup> Law of Thermodynamics

- A schematic representation of a heat engine. The engine receives energy  $Q_h$  from the hot reservoir, expels energy  $Q_c$  to the cold reservoir, and does work  $W$ .
- If working substance is a gas then we can use the PV diagram to track the process.

Physics 207: Lecture 28, Pg 9

### Heat Engines

- Example: The Stirling cycle

We can represent this cycle on a P-V diagram:

Physics 207: Lecture 28, Pg 10

### Heat Engines and the 2<sup>nd</sup> Law of Thermodynamics

- A heat engine goes through a cycle (start and stop at the same point, same state variables)
  - 1<sup>st</sup> Law gives  $\Delta U = Q + W = 0$
- What goes in must come out
  - 1<sup>st</sup> Law gives  $Q_h = Q_c + W_{\text{cycle}} \quad (Q\text{'s} > 0)$
- So (cycle mean net work on world)
  - $Q_{\text{net}} = |Q_h| - |Q_c| = -W_{\text{system}} = W_{\text{cycle}}$

Physics 207: Lecture 28, Pg 11

### Efficiency of a Heat Engine

- How can we define a "figure of merit" for a heat engine?
- Define the efficiency  $\epsilon$  as:

$$\epsilon = \frac{W_{\text{cycle}}}{|Q_h|} = \frac{|Q_h| - |Q_c|}{|Q_h|} = 1 - \frac{|Q_c|}{|Q_h|}$$

Observation: It is impossible to construct a heat engine that, operating in a cycle, produces no other effect than the absorption of energy from a reservoir and the performance of an equal amount of work

Physics 207: Lecture 28, Pg 12

# Physics 207 – Lecture 23

### Heat Engines and the 2<sup>nd</sup> Law of Thermodynamics

It is impossible to construct a heat engine that, operating in a cycle, produces no other effect than the absorption of energy from a reservoir and the performance of an equal amount of work.

This leads to the 2<sup>nd</sup> Law

Equivalently, heat flows from a high temperature reservoir to a low temperature reservoir

Physics 207: Lecture 28, Pg 13

### Lecture 28: Exercise 3 Efficiency

- Consider two heat engines:
  - Engine I:
    - Requires  $Q_{in} = 100$  J of heat added to system to get  $W=10$  J of work (done on world in cycle)
  - Engine II:
    - To get  $W=10$  J of work,  $Q_{out} = 100$  J of heat is exhausted to the environment
- Compare  $\epsilon_I$ , the efficiency of engine I, to  $\epsilon_{II}$ , the efficiency of engine II.
 
$$\epsilon = \frac{W_{cycle}}{|Q_h|} = \frac{|Q_h| - |Q_c|}{|Q_h|} = 1 - \frac{|Q_c|}{|Q_h|}$$

(A)  $\epsilon_I < \epsilon_{II}$       (B)  $\epsilon_I > \epsilon_{II}$       (C) Not enough data to determine

Physics 207: Lecture 28, Pg 14

### Reversible/irreversible processes and the best engine, ever

- Reversible process:
  - Every state along some path is an equilibrium state
  - The system can be returned to its initial conditions along the same path
- Irreversible process:
  - Process which is not reversible !
- All real physical processes are irreversible
  - e.g. energy is lost through friction and the initial conditions cannot be reached along the same path
  - However, some processes are almost reversible
    - If they occur slowly enough (so that system is almost in equilibrium)

Physics 207: Lecture 28, Pg 15

### The Carnot cycle

*Carnot Cycle*  
Named for Sadi Carnot (1796- 1832)

- 1) Isothermal expansion
- 2) Adiabatic expansion
- 3) Isothermal compression
- 4) Adiabatic compression

Physics 207: Lecture 28, Pg 16

### The Carnot Engine (the best you can do)

- No real engine operating between two energy reservoirs can be more efficient than a Carnot engine operating between the same two reservoirs.

- A → B, the gas expands isothermally while in contact with a reservoir at  $T_h$
- B → C, the gas expands adiabatically ( $Q=0, \Delta U=W_{B \rightarrow C}, T_h \rightarrow T_c$ ), PV<sup>γ</sup>=constant
- C → D, the gas is compressed isothermally while in contact with a reservoir at  $T_c$
- D → A, the gas compresses adiabatically ( $Q=0, \Delta U=W_{D \rightarrow A}, T_c \rightarrow T_h$ )

Physics 207: Lecture 28, Pg 17

### Carnot Cycle Efficiency

$$\epsilon_{Carnot} = 1 - Q_c/Q_h$$

$$Q_{A \rightarrow B} = Q_h = W_{AB} = nRT_h \ln(V_B/V_A)$$

$$Q_{C \rightarrow D} = Q_c = W_{CD} = nRT_c \ln(V_D/V_C)$$

(work done by gas)

But  $P_A V_A = P_B V_B = nRT_h$  and  $P_C V_C = P_D V_D = nRT_c$   
 so  $P_B/P_A = V_A/V_B$  and  $P_D/P_C = V_C/V_D$   
 as well as  $P_B V_B^\gamma = P_C V_C^\gamma$  and  $P_D V_D^\gamma = P_A V_A^\gamma$   
 with  $P_B V_B^\gamma / P_A V_A^\gamma = P_C V_C^\gamma / P_D V_D^\gamma$  thus

$$\rightarrow (V_B/V_A) = (V_D/V_C)$$

$$Q_c/Q_h = T_c/T_h$$

Finally

$$\epsilon_{Carnot} = 1 - T_c/T_h$$

Physics 207: Lecture 28, Pg 18

# Physics 207 – Lecture 23

### The Carnot Engine

- Carnot showed that the thermal efficiency of a Carnot engine is:

$$\epsilon_{\text{Carnot cycle}} = 1 - \frac{T_{\text{cold}}}{T_{\text{hot}}}$$

- All real engines are less efficient than the Carnot engine because they operate irreversibly due to the path and friction as they complete a cycle in a brief time period.

Physics 207: Lecture 28, Pg 19

### Power from ocean thermal gradients... oceans contain large amounts of energy

#### Carnot Cycle Efficiency

$$\epsilon_{\text{Carnot}} = 1 - Q_c/Q_h = 1 - T_c/T_h$$

See: <http://www.nrel.gov/otec/what.html>

Physics 207: Lecture 28, Pg 20

### Ocean Conversion Efficiency

$$\epsilon_{\text{Carnot}} = 1 - Q_c/Q_h = 1 - T_c/T_h$$

$$\epsilon_{\text{Carnot}} = 1 - T_c/T_h = 1 - 275 \text{ K}/300 \text{ K}$$

$$= 0.083 \text{ (even before internal losses and assuming a REAL cycle)}$$

Still: "This potential is estimated to be about  $10^{13}$  watts of base load power generation, according to some experts. The cold, deep seawater used in the OTEC process is also rich in nutrients, and it can be used to culture both marine organisms and plant life near the shore or on land."

"Energy conversion efficiencies as high as 97% were achieved."

See: <http://www.nrel.gov/otec/what.html>

So  $\epsilon = 1 - Q_c/Q_h$  always correct but  $\epsilon_{\text{Carnot}} = 1 - T_c/T_h$  only reflects a Carnot cycle

Physics 207: Lecture 28, Pg 21

### Lecture 28: Exercises 4 and 5

#### Free Expansion and the 2<sup>nd</sup> Law

You have an ideal gas in a box of volume  $V_1$ . Suddenly you remove the partition and the gas now occupies a large volume  $V_2$ .

- How much work was done by the system?
- What is the final temperature ( $T_2$ )?
- Can the partition be reinstated with all of the gas molecules back in  $V_1$ ?

1: (A)  $W > 0$                       (B)  $W = 0$                       (C)  $W < 0$

2: (A)  $T_2 > T_1$                       (B)  $T_2 = T_1$                       (C)  $T_2 < T_1$

Physics 207: Lecture 28, Pg 22

### Entropy and the 2<sup>nd</sup> Law

- Will the atoms go back?
  - Although possible, it is quite improbable
  - The are many more ways to distribute the atoms in the larger volume that the smaller one.
- Disorderly arrangements are much more probable than orderly ones
- Isolated systems tend toward greater disorder
  - Entropy ( $S$ ) is a measure of that disorder
  - Entropy ( $\Delta S$ ) increases in all natural processes. (The 2<sup>nd</sup> Law)
  - Entropy and temperature, as defined, guarantees the proper direction of heat flow.

all atoms      no atoms

Physics 207: Lecture 28, Pg 23

### Entropy and the 2<sup>nd</sup> Law

- In a reversible process the total entropy remains constant,  $\Delta S=0!$
- In a process involving heat transfer the change in entropy  $\Delta S$  between the starting and final state is given by the heat transferred  $Q$  divided by the absolute temperature  $T$  of the system.

$$\Delta S \equiv \frac{Q}{T}$$


- The 2<sup>nd</sup> Law of Thermodynamics
 

"There is a quantity known as entropy that in a closed system always remains the same (reversible) or increases (irreversible)."
- Entropy, when constructed from a microscopic model, is a measure of disorder in a system.

Physics 207: Lecture 28, Pg 24

# Physics 207 – Lecture 23


### Entropy, Temperature and Heat



- Example: Q joules transfer between two thermal reservoirs as shown below
- Compare the total change in entropy.

$$\Delta S = (-Q/T_1) + (+Q/T_2) > 0$$

because  $T_1 > T_2$



Physics 207: Lecture 28, Pg 25

### Entropy and Thermodynamic processes

Examples of Entropy Changes:

Assume a reversible change in volume and temperature of an ideal gas by expansion against a piston held at constant pressure ( $dU = dQ - P dV$  with  $PV = nRT$  and  $dU/dT = C_v$ ):

$$\Delta S = \int_i^f dQ/T = \int_i^f (dU + PdV) / T$$

$$\Delta S = \int_i^f \{C_v dT / T + nR(dV/V)\}$$

$$\Delta S = nC_v \ln (T_f / T_i) + nR \ln (V_f / V_i)$$


Ice melting:

$$\Delta S = \int_i^f dQ/T = Q/T_{\text{melting}} = m L_f / T_{\text{melting}}$$

Physics 207: Lecture 28, Pg 26

### The Laws of Thermodynamics

- First Law  
**You can't get something for nothing.**
- Second Law  
**You can't break even.**
- **Do not forget: Entropy, S, is a state variable**



Physics 207: Lecture 28, Pg 27

### Recap, Lecture 28

- Agenda: Ch. 21 Finish, Start Ch. 22
  - ❖ Ideal gas at the molecular level, Internal Energy
  - ❖ Molar Specific Heat ( $Q = m c \Delta T = n C \Delta T$ )
  - ❖ Ideal Gas Molar Heat Capacity (and  $\Delta U_{\text{int}} = Q + W$ )
    - Constant V:  $C_v = 3/2 R$ , Constant P:  $C_p = 3/2 R + R = 5/2 R$
  - ❖ Adiabatic processes (no external heat transfer)
  - ❖ Heat engines and Second Law of thermodynamics
  - ❖ Reversible/irreversible processes and Entropy

Assignments:

- Problem Set 10 (Ch. 20 & 21) due Tuesday, Dec. 12, 11:59 PM  
Ch. 20: 13,22,38,43,50,68 Ch.21: 2,16,29,36,70
- Problem Set 11, Ch. 22: 6, 7, 17, 37, 46 (Due, Friday, Dec. 15, 11:59 PM)
- Wednesday, Start Problem Set 11

Physics 207: Lecture 28, Pg 28