

Title: "Free" electrons in a periodic potential

Bloch's Theorem:

A solution of the Schrödinger equation in a periodic lattice (i.e., with a periodic potential) can be changed by only a phase factor for all lattice translations. In other words, the probability density must be the same for all equivalent positions in the lattice. Hence  $\psi^*(\vec{r}) \psi(\vec{r}) = \psi^*(\vec{r} + \vec{R}) \psi(\vec{r} + \vec{R})$  where  $\vec{R} \equiv n_1 \vec{a}_1 + n_2 \vec{a}_2 + n_3 \vec{a}_3$ .

In general:

$$\begin{aligned} \psi_k(\vec{r}) &= e^{i\vec{k}\cdot\vec{r}} u(\vec{r}) \quad \text{where } e^{i\vec{k}\cdot\vec{r}} \text{ is a plane wave} \\ u(\vec{r}) &\equiv \vec{u}(\vec{r} + \vec{R}) \quad \text{and is now assumed periodic} \\ \psi_k(\vec{r} + \vec{R}) &= e^{i\vec{k}\cdot(\vec{r} + \vec{R})} u(\vec{r} + \vec{R}) \\ \psi_k(\vec{r} + \vec{R}) &= e^{i\vec{k}\cdot\vec{R}} e^{i\vec{k}\cdot\vec{r}} u(\vec{r}) \\ \psi_k(\vec{r} + \vec{R}) &= e^{i\vec{k}\cdot\vec{R}} \psi_k(\vec{r}) \end{aligned}$$

Not only does  $\psi_k(\vec{r} + \vec{R}) = e^{i\vec{k}\cdot\vec{R}} \psi_k(\vec{r})$ , it also is multi-valued.

$$\begin{aligned} \text{Let } \vec{k} = \vec{G} + \vec{k}' \quad e^{i\vec{k}\cdot\vec{R}} &= e^{i(\vec{G} + \vec{k}')\cdot\vec{R}} = e^{i\vec{G}\cdot\vec{R}} e^{i\vec{k}'\cdot\vec{R}} \\ &= e^{i2\pi n} e^{i\vec{k}'\cdot\vec{R}} \end{aligned}$$

so

$$\psi_k(\vec{r} + \vec{R}) = e^{i\vec{k}'\cdot\vec{R}} \psi_k(\vec{r})$$

but we could have written equivalently

$$\psi_{k'}(\vec{r} + \vec{R}) = e^{i\vec{k}'\cdot\vec{R}} \psi_{k'}(\vec{r})$$

and so  $\psi_k$  and  $\psi_{k'}$  must have the same phase shift.

Thus a general  $\psi(\vec{r})$  solution must use a composite sum

$$\psi(\vec{r}) = \sum_{\vec{G}} \psi_{\vec{k}-\vec{G}}(\vec{r}) u_{\vec{k}-\vec{G}}(\vec{r})$$

Notice  $\vec{k}$  is the only wave vector on the right hand side

$$\psi_k(\vec{r}) = \sum_{\vec{G}} \psi_{\vec{k}-\vec{G}}(\vec{r}) u_{\vec{k}-\vec{G}}(\vec{r})$$

So we must define an additional parameter to specify which solution.

$$\psi_{n,\vec{k}} = e^{i\vec{k}\cdot\vec{r}} u_{n,\vec{k}}(\vec{r}) \quad n \equiv \text{band index.}$$

In the case of nearly free electrons  $U_0$  is assumed to be constant.

Using the time independent Schrödinger equation,  $-\frac{\hbar^2}{2m}\nabla^2\psi(\vec{r}) + U_0\psi(\vec{r}) = E\psi(\vec{r})$ , and letting  $\vec{G} = 0$  and  $\psi_k = A e^{i\vec{k}\cdot\vec{r}}$


$$\text{Normalization requires } \int_{x\text{-total}} \psi^* \psi dV = 1 \quad A = \frac{1}{\sqrt{V}}$$

with energy eigenvalue,  $E = U_0 + \frac{\hbar^2 k^2}{2m}$ , and eigenfunction,  $\psi_k = \frac{1}{\sqrt{V}} e^{i\vec{k}\cdot\vec{r}}$ .

Notice that we could choose another solution  $\vec{k} \Rightarrow \vec{k} + \vec{G}$

$$\text{This gives } \psi = A e^{i(\vec{k}+\vec{G})\cdot\vec{r}} \text{ and } \nabla^2\psi = -A|\vec{k} + \vec{G}|^2\psi$$

so that  $E = U_0 + \frac{\hbar^2}{2m}|\vec{k} + \vec{G}|^2$



Reduced Zone: Look at all solutions in the first Brillouin zone

Extended Zone: Look at solutions in the appropriate zone

So what happens at the zone boundary?

Bragg reflection of the electrons!

We might therefore expect equal numbers of the two possible solutions.

$$A e^{i\frac{\vec{G}}{2}\cdot\vec{x}} \quad \text{and} \quad A e^{-i\frac{\vec{G}}{2}\cdot\vec{x}} \quad \text{where} \quad |G| = \frac{2\pi}{a}$$

$$\psi(+)=\exp\left(i\frac{G}{2}x\right)+\exp\left(-i\frac{G}{2}x\right)=2\cos(\pi x/a)$$

and

$$\psi(-)=\exp\left(i\frac{G}{2}x\right)-\exp\left(-i\frac{G}{2}x\right)=2i\sin(\pi x/a)$$

$$\psi^*(+)\psi(+)\propto\cos^2(\pi x/a)$$


$$\psi^*(-)\psi(-)\propto\sin^2(\pi x/a)$$

The + solution builds up charge at ions

while the - solution builds up charge between ions

At present  $U$  is constant and thus there is no difference in the eigenvalues


If  $U$  is not constant (generally attractive), then




$|\psi(+)|^2$  is lower in energy

$|\psi(-)|^2$  is higher in energy

There are gaps wherever “bands” cross.



If we have a single electron and apply an electric field: (No scattering)



Magnitude of the energy gap

$$E_g = E_+ - E_-$$

$$E_+ = \int_0^1 dx \psi^*(+) U(x) \psi(+)$$

$$E_- = \int_0^1 dx \psi^*(-) U(x) \psi(-)$$

if  $U(x) = U_0 \cos\left(\frac{2\pi x}{a}\right)$  with period  $a$ !

$$E_{\text{gap}} = 2 \int dx U_0 \cos\left(\frac{2\pi x}{a}\right) \left[ \cos^2\left(\frac{\pi x}{a}\right) - \sin^2\left(\frac{\pi x}{a}\right) \right]$$

(between 1st and 2nd bands)

We can always decompose  $U$  in terms of its *Fourier* components to determine the gap at each and every band crossing.